

The O-Na and Mg-Al Anticorrelations in Turn-Off and early Subgiants in Globular Clusters^{*}

R.G. Gratton¹, P. Bonifacio², A. Bragaglia³, E. Carretta¹, V. Castellani⁴, M.
Centurion², A. Chieffi⁵, R. Claudi¹, G. Clementini³, F. D'Antona⁶, S.
Desidera^{1,7}, P. François⁸, F. Grundahl⁹, S. Lucatello^{1,7}, P. Molaro², L.
Pasquini⁸, C. Sneden¹⁰, F. Spite¹¹, and O. Straniero¹²

¹ Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, 35122 Padova,
Italy

² Osservatorio Astronomico di Trieste, Italy

³ Osservatorio Astronomico di Bologna, via Ranzani 1, 40127 Bologna, Italy

⁴ Dipartimento di Fisica, Università di Pisa, Italy

⁵ Istituto di Astrofisica Spaziale, CNR, Italy

⁶ Osservatorio Astronomico di Roma, Italy

⁷ Dipartimento di Astronomia, Università di Padova, Italy

⁸ European Southern Observatory

⁹ Department of Astronomy, University of Aarhus, Denmark

¹⁰ The University of Texas at Austin, USA

¹¹ Observatoire de Meudon, France

¹² Osservatorio Astronomico di Collurania, Italy

Received ; accepted

^{*} Based on data collected at the European Southern Observatory, Chile

Abstract. High dispersion spectra ($R \gtrsim 40,000$) for a quite large number of stars at the main sequence turn-off and at the base of the subgiant branch in NGC6397 and NGC6752 were obtained with the UVES on Kueyen (VLT UT2). The $[\text{Fe}/\text{H}]$ values we found are $-2.03 \pm 0.02 \pm 0.04$ and $-1.42 \pm 0.02 \pm 0.04$ for NGC6397 and NGC6752 respectively, where the first error bars refer to internal and the second ones to systematic errors. In both clusters the $[\text{Fe}/\text{H}]$'s obtained for TO-stars agree perfectly (within a few per cents) with that obtained for stars at the base of the RGB. The $[\text{O}/\text{Fe}] = 0.21 \pm 0.05$ value we obtain for NGC6397 is quite low, but it conforms to previous results obtained for giants in this cluster. Moreover, the star-to-star scatter in both O and Fe is very small, indicating that this small mass cluster is chemically very homogenous. On the other side, our results show clearly and for the first time that the O-Na anticorrelation (up to now seen only for stars on the red giant branches of globular clusters) is present among unevolved stars in the globular cluster NGC6752, a more massive cluster than NGC6397. A similar anticorrelation is present also for Mg and Al, and C and N. It is very difficult to explain the observed Na-O, and Mg-Al anticorrelation in NGC6752 stars by a deep mixing scenario; we think it requires some non internal mechanism.

Key words. Stars: abundances – Stars: evolution – Stars: Population II – Galaxy: globular clusters

1. INTRODUCTION

Globular clusters (GCs) are (with a few significant exceptions, most notably the case of ω Cen) generally very homogenous as far as Fe-peak elements are considered; on the other side, abundances of the lighter elements (from C to Al) show a complex, not yet adequately explained pattern. Rather wide variations in the strength of the CH and CN bands among stars on the red giant branch (RGB) were already noticed in the seventies (Osborn 1971, and many other references; for reviews see Smith 1987, Kraft 1994), and found to be correlated with other cluster properties (Norris 1987). More recently, a CN-CH anticorrelation was found in Main Sequence (MS) stars of NGC 6752 (Suntzeff & Smith 1991), 47 Tuc (Cannon et al. 1998), and M 71 (Cohen 1999a) from low dispersion spectra. Even more striking is the Na-O anticorrelation among GC red giants discovered by the Lick-Texas group (see e.g. Kraft et al. 1997).

These features are indications of the presence of elements processed through the complete CNO-cycle in the atmospheres of some globular cluster stars; at temperatures where complete CNO cycle occurs, quite large amounts of Na may be produced by proton capture on ^{22}Ne nuclei (Denissenkov & Denissenkova 1990; Langer & Hoffman 1995; Cavallo, Sweigart & Bell 1996). Whatever the cause of these abundance anomalies, it appears to be related to the dense environment of GCs. In fact, field halo stars present

surface abundances of Li, C, N, O, Na that are well explained by standard models, once a second mixing episode (only affecting Li, C and N abundances) after the RGB-bump is allowed (Gratton et al. 2000). Note that the RGB bump plays an important rôle, because there the molecular weight barrier created by the deepest inward penetration of the outer convective envelope (at the base of the RGB) is canceled by the outward shift of the H-burning shell. The absence of this barrier in the following evolution allows deep mixing caused by e.g. meridional circulations (Sweigart & Mengel 1979; Charbonnel 1994; Denissenkov & Tout 2000). Hence, both the O-Na anticorrelation observed among giants, and the C and N anomalies observed in stars in different evolutionary phases, are peculiar to GCs.

The proposed explanations of the abundance anomalies in globular cluster stars have been oscillating in years between an "in situ" mechanism (very deep mixing in RGB stars, possibly activated by a core rotation that might be larger in cluster than in field stars) and an external source of material (either primordial proto-cluster gas or processed material from a polluting companion). Since MS stars do not have a convective envelope able to enhance N while depleting C, these anomalies should be attributed mainly to a primordial origin and/or to pollution, while evolutionary mixing could possibly be responsible only for likely minor effects in RGB stars. However, rather surprisingly, an analysis of stars below the Turn-off (TO) of M 13 (Cohen 1999b) shows no variations in the strength of CN and CH molecular features, likely excluding primordial variation of C and N abundance in this cluster, which instead presents the clearest Na-O anticorrelation on the RGB (Kraft et al. 1997), and a range in the strength of CN and CH bands among giants (Suntzeff 1981). While the result of Cohen is very surprising, the analysis is only qualitative (a quantitative analysis being planned in a future, still unpublished paper), so that it is not clear which is the upper limit found to C and N abundance variations among MS stars in this very important cluster. In general, the relation between the Na-O and C-N anticorrelation is not yet well established. Observation of a Na-O anticorrelation among main sequence stars of globular clusters would be crucial, clearly ruling out the deep mixing scenario.

Earlier explorative studies of stars close to the TO and at the base of the subgiant branch of a few clusters have been presented by King, Stephens & Boesgaard (1998), Deliyannis, Boesgaard & King (1995) and Boesgaard et al. (1998) who used HIRES at Keck to gather spectra of three stars in M92; and Pasquini & Molaro (1996, 1997) who exploited the presence of closer clusters in the southern hemisphere to obtain spectra of a few stars in NGC6397 and 47 Tuc using EMMI at NTT (these stars are about a magnitude brighter than those of M92). However, only a handful of stars could be observed, and at very low S/N. Only UVES (D'Odorico et al. 2000) at the Kueyen (VLT UT2) Telescope is able to provide abundances of O, Na, Mg, Al and other elements for significant samples of well studied stars at the TO and at the base of RGB (well below the RGB bump),




Fig. 1. Positions of program stars (large symbols) on the Strömgren colour-magnitude diagram of NGC6397 (panel a) and NGC6752 (panel b). Photometry is from Grundahl et al. (1999). Note that photometry for NGC6397 is still uncalibrated, so that only instrumental magnitudes are given

crucial to clarify most of these issues. Here we present a short description of the first results of the ESO Large Program 165.L-0263 devoted to acquisition and analysis of high dispersion spectra ($R \gtrsim 40,000$) of a rather large sample of stars at the TO and at the base of the RGB in NGC6752, NGC6397, and 47 Tuc. A more detailed presentation of the analysis methods, and of related results for Li, will be given in forthcoming papers, still in preparation.

2. SAMPLE SELECTION AND OBSERVATIONS

Observations were carried out in two 6-nights runs (June and September 2000); due to poor weather conditions in the September run, we were not able to complete the acquisition of the spectra in 47 Tuc. Hence the present paper deals only with data for the




Fig. 2. Colour-magnitude diagram for the field stars; M_V is computed using Hipparcos parallaxes

two other clusters. In the same runs we also acquired data for 25 field stars with good parallaxes from the Hipparcos catalogue and metallicity similar to that of the GCs.

For each cluster, we selected two groups of stars: (i) dwarfs about 0.2 mag brighter than the turn-off (5 stars in NGC6397 and 9 in NGC6752); and (ii) stars at the base of the red giant branch (3 stars in NGC6397 and 9 stars in NGC6752), hereinafter called subgiants. Positions of the observed stars in the c-m diagrams of the two clusters (from Grundahl et al. 1999) are given in Figure 1. A posteriori, radial velocities and chemical composition show that all stars are bona fide members of the clusters. Stars with likely strong and weak CN bands were selected from Strömgren photometry: the c_1 index seems in fact to be higher for CN-rich stars (see Grundahl et al. 1999). Given this choice, it should be clear that stars are not a random sample of stars in the different magnitude bins, but rather they were selected in order to include quite extreme cases of CN-poor and CN-rich stars. For comparison, we give in Figure 2 also the location in a similar c-m diagram of the field stars we observed.

The observations were carried out using the dichroic beamsplitter #2. This allowed to observe simultaneously two spectral ranges with the two arms of the spectrograph. Slightly different setups were used in the two runs: the blue spectra cover the spectral range from about 3,500 Å to about 4,700 Å, while the red spectra cover the range from 5,700 Å to about 8,700 Å. This very broad spectral range allows observations of a large number of spectral features. The resolution was set by the slit width: in most cases this was 1 arcsec (yielding a resolution of 43,000 at order centers); however, this value

Table 1. Parameters for field stars

Star	M_v	$B - V$	$b - y$	[Fe/H]	$\log g$	T_{eff}	T_{eff}	T_{eff}
HD/BD						H $_{\alpha}$	Best	Alonso et al.
10607	4.03	0.56	0.372	-0.95	4.00	5780	5734	
29907	6.01	0.63	0.452	-1.47	4.82	5636	5406	
31128	5.05	0.49	0.358	-1.44	4.63	6279	5969	
108177	4.87	0.43	0.330	-1.70	4.44	5884	6063	6067
116064	4.74	0.45	0.349	-1.87	4.35	5770	5923	
120559	5.93	0.66	0.424	-0.90	4.51	5207	5378	
121004	5.12	0.62	0.424	-0.75	4.32	5401	5548	
126681	5.71	0.60	0.400	-1.13	4.55	5444	5556	5541
132475	3.62	0.56	0.401	-1.61	3.79	5596	5520	5788
134169	3.82	0.565	0.370	-0.81	4.06	6106	5825	5874
134439	6.72	0.77	0.487	-1.31	4.78	5151	5052	4974
134440	7.05	0.85	0.527	-1.28	4.69	4777	4777	4746
140283	3.34	0.49	0.380	-2.46	3.67	5560	5645	5691
145417	6.81	0.82	0.509	-1.23	4.78	5096	4922	
159482	4.92	0.58	0.382	-0.83	4.32	5546	5656	
166913	4.22	0.45	0.326	-1.59	4.08	5921	6020	
181743	4.96	0.45	0.351	-1.81	4.42	5749	5927	
188510	5.59	0.60	0.416	-1.44	4.50	5412	5485	5564
189558	3.59	0.575	0.386	-1.15	3.87	5829	5670	5663
193901	5.44	0.55	0.377	-1.03	4.62	5848	5780	5750
194598	4.60	0.49	0.348	-1.15	4.42	6064	5980	6017
204155	4.07	0.57	0.378	-0.70	4.06	5907	5751	
205650	5.39	0.51	0.375	-1.12	4.52	5628	5782	
-35 0360	6.31	0.76	0.472	-1.08	4.52	4957	5086	4980
+05 3640	6.50	0.74	0.474	-1.15	4.62	4987	5063	

was sometimes changed according to the variable seeing conditions, from 0.7 arcsec in the best cases, down to 1.2 arcsec in the worst ones. The slit length was always set at 8 arcsec, allowing accurate sky subtraction (necessary, since observations were carried out in full moon). Most of the objects are very faint ($V \gtrsim 17$), so that long exposures were required even with an 8 m telescope like Kueyen, and a very efficient spectrograph like UVES. Typical exposure times were ~ 1 hr for subgiants, and about 4 hr (splitted in 3 to 4 segments) for TO stars. The S/N (per pixel at 6700 Å; there are about 5 pixels per resolution element) of the resulting spectra is ~ 100 in the best cases, and down to about 20 in the worst ones. Relevant data are presented in Tables 1, 2, and 3.

Table 2. Data for NGC 6397 stars (identifications from Grundahl et al. 1999); Al abundances from resonance lines

Star	S/N	n.	[Fe/H]	r.m.s	EW	[O/Fe]	[O/Fe]	EW	EW	[Na/Fe]	[Mg/Fe]	[Al/Fe]
		lines			7771	LTE	nLTE	8184	8190	nLTE		
Dwarfs												
1543	91	23	-2.02	0.12	14.7	0.32	0.16	..	38.8	0.21	0.06	-0.40
1622	82	21	-2.02	0.10	13.3	0.27	0.11	28.5	40.7	0.28	0.10	-0.34
1905	92	22	-2.06	0.11	13.4	0.27	0.11	17.1	29.4	0.02	0.12	-0.35
201432	97	22	-2.00	0.11	12.8	0.25	0.08	17.0	35.6	0.08	0.06	-0.50
202765	59	18	-2.02	0.14	16.1	0.38	0.21	20.7	28.5	0.06	-0.06	-0.41
Subgiants												
669	91	27	-2.01	0.11	6.0	0.34	0.26	62.9	86.8	0.48	0.23	-0.06
793	105	31	-2.04	0.12	<6.0	<0.34	<0.26	39.9	72.1	0.21	0.28	-0.21
206810	85	23	-2.10	0.10	8.0	0.48	0.48	49.5	68.3	0.25	0.10	-0.24

Table 3. Data for NGC 6752 stars. Stars are ordered according to increasing Na abundance. Star identifications are from Grundahl et al. (1999). Note: 1) Al abundances from resonance lines; (2) Al abundances from high excitation 8772.9/73.9 doublet

Star	S/N	n.	[Fe/H]	r.m.s	EW	[O/Fe]	[O/Fe]	EW	EW	[Na/Fe]	[Mg/Fe]	[Al/Fe]
		lines			7771	LTE	nLTE	8184	8190	nLTE		
Dwarfs												
4428	49	22	-1.52	0.14	35.7	0.35	0.24	24.4	57.1	-0.35	0.04	-1.07 (1)
4383	49	23	-1.42	0.19	46.0	0.60	0.49	33.6	61.5	-0.23	0.01	-0.70 (1)
202316	44	21	-1.56	0.20	28.7	0.31	0.20	42.6	69.0	-0.09	-0.06	-1.04 (1)
4341	42	23	-1.57	0.26	21.6	0.30	0.21	63.4	83.7	0.18	0.06	-1.35 (1)
4458	33	17	-1.52	0.27	19.4	0.04	-0.06	71.9	82.6	0.24	-0.09	-1.22 (1)
4661	21	16	-1.24	0.17	..	..	..	75.4	84.3	0.28	0.15	-0.68 (1)
5048	44	25	-1.43	0.12	11.0	-0.29	-0.37	78.2	93.7	0.37	-0.24	-0.40 (1)
4907	66	28	-1.41	0.21	12.0	-0.25	-0.34	102.9	103.0	0.61	-0.09	-0.45 (1)
200613	25	18	-1.24	0.18	..	..	..	103.5	110.1	0.64	0.06	0.01 (1)
Subgiants												
1406	48	44	-1.39	0.16	18.0	0.40	0.34	82.1	105.7	0.02	0.20	0.10 (2)
1665	49	41	-1.34	0.15	..	..	..	86.7	116.6	0.10	0.10	-0.13 (2)
1445	35	33	-1.53	0.13	..	..	..	95.0	..	0.19	0.06	0.33 (2)
1563	44	38	-1.42	0.17	17.4	0.38	0.33	97.0	131.0	0.25	0.11	0.17 (2)
1400	70	41	-1.44	0.14	15.3	0.33	0.28	103.4	..	0.29	0.10	0.34 (2)
1461	25	32	-1.38	0.24	..	..	..	94.9	144.4	0.29	0.10	-0.06 (2)
202063	29	38	-1.49	0.27	..	..	..	120.4	143.8	0.44	0.13	0.33 (2)
1481	55	46	-1.39	0.15	..	..	..	122.5	166.2	0.54	-0.07	0.69 (2)
1460	38	32	-1.37	0.15	<14.0	<0.25	<0.19	137.4	..	0.65	-0.19	0.86 (2)

3. ANALYSIS

Equivalent widths (EWs) for a large number of lines were measured using an automatic procedure (similar to the one used in Bragaglia et al. 2001; full details will be given elsewhere). Typical errors in the EWs are ± 3 mÅ for NGC6397 stars, and ± 5 mÅ for NGC6752 stars, but they are as low as ± 2.5 mÅ in the best cases (the Li lines at 6707 Å). These values are comparable with results obtained for much brighter stars at other telescopes.

One of the aims of the present analysis was to derive accurate reddening estimates for GCs by comparing the observed colours with effective temperatures T_{eff} 's obtained from the spectra; for this reason our temperatures are based on fitting of the wings of the H_α profiles with synthesized lines¹. The model lines were computed according to the precepts given in Castelli, Gratton & Kurucz (1997); throughout this paper we used model atmospheres extract by interpolation within the grid by Kurucz with the overshooting option switched off (Kurucz 1993). Within these models, convection is considered using a mixing length approach, with a pressure scale height value of $l/H_p = 1.25$. Internal errors of individual temperatures are estimated to be 150-200 K for individual stars; these rather large errors are mainly due to uncertainties in the flat fielding procedure, and are typical for such temperature derivations when only one Balmer line is used (see e.g. Fuhrmann, Axer & Gehren 1994). The adopted temperature scale is confirmed by analysis of line excitation: however, temperature from excitation have slightly larger error bars due to the limited range of excitation for lines measurable in warm, metal-poor dwarfs and subgiants.

For field stars, we compared these T_{eff} 's (Column 7 of Table 1) with those given by colours ($B-V$ and $b-y$), using the calibration given by Kurucz (1993); here we assumed that all these nearby (field) stars are unreddened. Internal errors in T_{eff} 's from colours are about 26 K. On average, temperatures derived from H_α are lower than those derived from colours by 97 ± 38 K, with a large r.m.s scatter of 188 K. For field stars, we then corrected the temperatures from colours to those derived from H_α , and averaged the two values, giving a weight 4 to temperatures from colours, and 1 to temperatures derived from H_α . These are our "best temperatures" for the field stars (Column 8 of Table 1).

We may compare these "best temperatures" with those obtained by Alonso, Arribas & Martinez-Roger (1996) using the IR flux method (Column 9 of Table 1). Excluding the discrepant case of HD132475, on average our best T_{eff} 's are lower than those given by Alonso et al. by 5 ± 14 K (r.m.s. residuals=45 K, 10 stars). Hence our T_{eff} 's can be assumed to be on the same scale of Alonso et al.

¹ We only used H_α . H_β was not observed; the other members of the Balmer series were indeed observed, but they are more contaminated by metal lines, and their dependence on gravity, metal abundance and details in the convection much larger than for H_α (see Fuhrmann et al. 1993)

Table 4. Sensitivities of abundances to errors in the atmospheric parameters

Element	ΔT_{eff}	$\Delta \log g$	Δv_t
	+100 K	+0.2 dex	+0.2 km s ⁻¹
TO-star			
[Fe/H]	+0.096	−0.054	−0.050
[O/Fe]	−0.161	+0.123	+0.048
[Na/Fe]	−0.037	+0.007	+0.032
[Mg/Fe]	−0.034	+0.006	+0.041
[Al/Fe]	+0.001	−0.012	+0.007
Subgiants			
[Fe/H]	+0.108	−0.023	−0.049
[O/Fe]	−0.200	+0.098	+0.044
[Na/Fe]	−0.044	−0.010	+0.026
[Mg/Fe]	−0.055	+0.020	+0.043
[Al/Fe]	−0.080	+0.021	+0.038

When plotted one on top of the other (separately: dwarfs and subgiants, and stars in different clusters), the H_α profiles for cluster stars look undistinguishable. Furthermore, we were not able to find any correlation between colours and temperatures either from line excitation or H_α profiles. We then concluded that the stars are intrinsically very similar, and that the slightly different values of colours and temperatures we derived for each star are due to random errors. In the following analysis, we have then adopted for all stars in these groups the same average temperatures (note however that this assumption is not critical in the present discussion).

Surface gravities $\log g$ were obtained from the location of the stars in the colour magnitude diagram, assuming masses consistent with an age of 14 Gyr (again, this assumption is not critical). As usual, microturbulent velocities were obtained by eliminating trends of abundances derived from individual Fe lines with expected line strength. Star-to-star scatter in Fe abundances within each group (same cluster, same evolutionary phase) were reduced by adopting for all stars the same microturbulent velocity; these average values were then adopted in the final analysis.

The finally adopted atmospheric parameters (effective temperatures in K/surface gravities/model metal abundances/microturbulent velocities in km s⁻¹) were as follows: NGC6397 TO-stars (6476/4.10/−2.04/1.32); NGC6397 subgiants (5478/3.42/−2.04/1.32); NGC6752 TO-stars (6226/4.28/−1.43/0.70); NGC6752 subgiants (5347/3.54/−1.43/1.10).

Sensitivities of abundances to errors in the atmospheric parameters are given in Table 4. Errors in final abundances are mostly due to possible errors in the adopted T_{eff} 's. For our line list (dominated by low excitation lines), Fe I abundances raise by

0.10 dex, $[\text{O}/\text{Fe}]$ values by -0.20 dex, and $[\text{Na}/\text{Fe}]$ ones by -0.04 dex for a 100 K increase in the adopted T_{eff} . Systematic errors in T_{eff} 's are dominated by uncertainties in the fitting of the H_α profiles: they are about ± 90 K for the dwarfs and ± 60 K for the subgiants, leading to errors in the $[\text{Fe}/\text{H}]$ values of ± 0.09 dex and ± 0.06 dex respectively for TO-stars and subgiants. This value, appropriate for cluster stars, is smaller than that given for field stars simply because our T_{eff} 's for cluster stars are actually the average over the values obtained for several stars, and errors for individual stars are given by uncertainties in the flat fielding procedure, that are only weakly affected by S/N. Corresponding errors in $[\text{O}/\text{Fe}]$'s are ± 0.18 and ± 0.12 dex; those in $[\text{Na}/\text{Fe}]$ are ± 0.04 and ± 0.03 dex. However, in the context of the O-Na anticorrelation, errors in temperatures adopted for individual stars are more important. In the case of NGC6397, a quite realistic estimate can be obtained by the star-to-star scatter in Fe abundances, that is 0.032 dex, corresponding to an r.m.s. spread of 33 K in the T_{eff} 's. The adoption of a uniform temperature would then yield errors of ± 0.07 dex in $[\text{O}/\text{Fe}]$'s and ± 0.01 dex in $[\text{Na}/\text{Fe}]$'s (to be compared with the observed star-to-star scatter). In the case of NGC6752, the star-to-star scatter of 0.096 dex in $[\text{Fe}/\text{H}]$'s also includes an important contribution due to noise in the *EW*s: in fact the r.m.s. reduces to 0.074 dex if only spectra with $S/N > 40$ are considered. We conclude that errors in individual T_{eff} 's are $\lesssim 73$ K. Related errors in $[\text{O}/\text{Fe}]$ and $[\text{Na}/\text{Fe}]$ abundances are $\lesssim 0.15$ and $\lesssim 0.03$ dex. Again, these are the values to be compared with the observed scatter.

4. IRON ABUNDANCES

Fe abundances were derived using a rather large number of lines, typically ~ 20 and $\sim 30\text{--}40$ for TO-stars and subgiants respectively. Full details about the line list, including *gf*'s, *EW*s and abundances from individual lines, will be given in separate papers (in preparation).

The Fe abundance obtained for dwarfs in NGC6397 ($[\text{Fe}/\text{H}] = -2.02 \pm 0.01$) is in good agreement with that determined from subgiants ($[\text{Fe}/\text{H}] = -2.05 \pm 0.03$). The average of the two groups is $[\text{Fe}/\text{H}] = -2.03 \pm 0.02$ (this is the internal error of our analysis; systematic errors are ± 0.04 dex). This value is less than that derived from giants in Carretta & Gratton (1997: $[\text{Fe}/\text{H}] = -1.82 \pm 0.04$), and Zinn & West (1984: $[\text{Fe}/\text{H}] = -1.91$). However it agrees very well with the value obtained by Minniti et al. (1993: $[\text{Fe}/\text{H}] = -1.96 \pm 0.04$), and the recent, comprehensive analysis of giants and subgiants by Castilho et al. (2000: -2.00 ± 0.05).

The equilibrium of ionization is not well reproduced: abundances from neutral Fe lines are 0.11 dex larger than those from singly ionized Fe lines. The same result is obtained for NGC6752 and our field stars. Note that a smaller difference (0.07 dex) in the same sense is also present in our solar reference analysis (in that case we obtained $\log n(\text{Fe}) = 7.52$

from Fe I lines, and 7.45 from Fe II lines using the solar model atmosphere from Kurucz 1993, with no overshooting). We then think most of the difference for the Sun is due to either the adopted $gf's$ values (these are laboratory values taken from recent literature) or to the model atmospheres (models might underestimate the temperature gradient in real atmospheres, perhaps due to an inappropriate consideration of convection). The residual difference for the program stars might be due to the adoption of slightly too high T_{eff} 's ($\sim 40 - 50$ K) or too low gravities (by ~ 0.1 dex). Note that it cannot be due to departures from LTE, because the expected dominating effect (overionization: Idiart & Thévenin 1999; Gratton et al. 1999) would lead to larger abundances from Fe II lines than from Fe I ones (opposite to observations).

The star-to-star scatter in $[\text{Fe}/\text{H}]$ values is extremely small: the r.m.s. scatter is only 0.04 dex (i.e. 10%) for NGC 6397. This seems a very homogeneous cluster as far as Fe abundances are considered.

On the other side, the $[\text{Fe}/\text{H}]$ value for NGC6752 ($[\text{Fe}/\text{H}] = -1.42 \pm 0.02$, internal error; systematic error is again ± 0.04 dex), obtained both from TO and subgiant stars, which agree completely, coincides with what derived from giants by Carretta & Gratton (1997: $[\text{Fe}/\text{H}] = -1.42 \pm 0.02$); it is somewhat larger than the value quoted by Zinn & West ($[\text{Fe}/\text{H}] = -1.54$) and Minniti et al. (1993: $[\text{Fe}/\text{H}] = -1.58 \pm 0.04$). The spectra of NGC6752 have a S/N lower than those in NGC6397, since we chose to observe more stars, even at a lower S/N. The scatter of abundances for individual lines (from 0.12 to 0.27 dex) is larger than that obtained for stars in NGC6397, roughly in agreement with the lower S/N.

For field stars we may compare the present Fe abundances with those derived by Carretta, Gratton & Sneden (2000). Limiting ourselves to only those stars for which Carretta et al. considered high dispersion abundances, the present Fe abundances are smaller on average by -0.05 ± 0.02 dex (11 stars, r.m.s.=0.08 dex). The slightly lower metal abundances are due to lower T_{eff} 's adopted in the present paper.

5. THE O-NA AND MG-AL ANTICORRELATION

In our analysis, Oxygen abundances were derived from the permitted infrared triplet (7771-74 Å): the forbidden lines are too weak in the program stars to be reliably measured, and we did not observe the region of the OH band in the near UV because the required exposure times would be prohibitive even for an 8 meter telescope. Na abundances are based on the quite strong doublet at 8184-90 Å, clearly visible in all stars. We checked that telluric lines were not blended with the stellar features. For both O and Na, our abundances included non-LTE corrections, computed following the precepts of Gratton et al. (1999): however, these corrections are small and they do not affect any of the conclusion reached in this paper. Mg abundances are based on a few high excitation lines

in the blue and yellow portion of the spectrum (typically three to four lines were measured for each star). Finally, whenever possible Al abundances were measured using the high excitation IR doublet at 8772.9/73.9 Å. However, this was possible only for subgiants in NGC6752, since in the other cases this doublet is too weak, and the only chance to measure Al abundances is by using the resonance doublet at 3944/61 Å. However, this lines, while stronger, are not ideal abundance indicators due to saturation and the presence of large departures from LTE (see e.g. François 1986; Beyley & Cottrell 1987; and the discussion in Gratton & Sneden 1988). We have not prepared yet an adequate model for the Al atom to account for such departures from LTE; anyway abundances obtained by an LTE analysis of these lines may still be useful to discuss the Mg-Al anticorrelation insofar only stars with very similar atmospheric parameters are considered, because in this case we expect that departures from LTE should be essentially the same for all such stars.

For NGC6397, the O abundance is $[O/Fe]=+0.21 \pm 0.05$ (internal error; systematic errors are ~ 0.1 dex), and the average $[Na/Fe]$ ratio is $[Na/Fe]=+0.20 \pm 0.05$. This O excess is quite small in comparison to the values usually found for metal-poor stars (see e.g. Gratton et al. 2000), but it agrees very well with the mean values determined by Minniti et al. (1997: $[O/Fe]=+0.19$) and Norris & Da Costa (1995: $[O/Fe] \sim 0.1$) from analysis of the forbidden lines in red giant spectra. Also the Mg abundance we get is quite low: this point will be discussed in a next paper. On the other side, the $[Na/Fe]$ value we found agrees well with that determined for a single red giant by Carretta (1994: $[Na/Fe]=+0.22$), and for two more by Norris & Da Costa (1995: $[Na/Fe] \sim 0.2$), while it is somewhat larger than the value of $[Na/Fe]=-0.01$ found by Minniti et al. (1997). We wish to remark that none of the program stars seem to be oxygen-poor and sodium-rich. The star-to-star scatter in our determinations (0.15 and 0.14 dex, r.m.s., respectively) is larger than expected from errors in the EWs alone (we would expect r.m.s. values of ~ 0.10 dex): however it may be justified by some star-to-star scatter in the atmospheric parameters (within the measuring errors), not accounted for in our analysis (we are assuming that all dwarfs and subgiants may be analyzed using the same model atmospheres), and by a small offset in the results for dwarfs and subgiants (again, likely due to small errors in the adopted set of atmospheric parameters, within our quoted uncertainties). Given the small spread in Fe abundances, and the absence of a clear O-Na anticorrelation, we conclude that NGC6397 is indeed a very homogenous cluster. On the whole, our results indicate that stars in this cluster conforms to the paradigm set by field stars, confirming earlier findings by Bell & Dickens (1980).

NGC6752 presents a very different scenario (this is not too surprising, in view of the variations of the strength of CN and CH bands among stars on the MS already noticed by Suntzeff & Smith 1991). There is a clear anti-correlation between O and Na abundances even for stars at the TO. This is clearly illustrated by Figures 3 and 4,




Fig. 3. Spectral region including the 8184-90 Å Na I doublet in NGC6752 TO-stars (stars are ordered according to decreasing Na abundances). The position of the Na lines is marked. Note that all these stars essentially have the same temperature, gravity, overall metal abundance and microturbulent velocity, so that observed variations in the line strengths can be directly interpreted as spread in the abundances

where we plotted the spectral regions including the Na and O lines in these stars (stars are ordered according to decreasing Na abundances). Notice that all these stars have the same (or at least very similar) temperature, gravity, overall metal abundance and microturbulent velocity, so that observed variations in the line strengths may be directly interpreted as spread in the abundances: then, in spite of the fact that a few of the spectra are somewhat noisy, this result does not depend on details of the analysis, but rather is a solid purely observational evidence, largely independent of all the assumptions made. We found that Na abundances span a range of almost an order of magnitude. O abundances also change by a rather large factor; while not as extreme as in M13, we found a rather extensive O-Na anticorrelation in NGC6752 too. This anti-correlation is shown in Figure 5. There is a small offset in the O-Na anticorrelation between TO-stars




Fig. 4. Same as Figure 3, but for the region including the 7711-74 Å OI triplet in NGC6752 TO-stars (stars are ordered according to decreasing Na abundances). The position of the O lines is marked.

and subgiants, but we think this is an artefact of the analysis (slightly incorrect offsets in temperatures and gravities).

There is some correlation between c_1 and the O-Na trends; hence, as expected, this can be used to select candidates. Note however that the scatter in this relation is quite large (see Figure 6).

Figure 7 displays results for Mg and Al in NGC6752. The star-to-star variations in the Al abundances are large: ~ 1 dex among subgiants, and even more among TO-stars. These large variations cannot be explained by observational errors, and (for TO-stars) by any plausible differential non-LTE effect among stars with very similar atmospheric parameters. On the other side, star-to-star variations in the Mg abundances are much smaller ($\lesssim 0.4$ dex), and they may be seen clearly only among subgiants; for TO-stars, the used Mg lines are quite weak and a larger scatter in the abundances is due to errors in the EW s. Data for subgiants, for which reliable Al abundances could be obtained from the high excitation Al I doublet at 8772.9/73.9 Å, indicate that also for these elements




Fig. 5. Run of the abundances of Na and of O for stars in NGC6752. Filled squares are TO-stars, open squares are subgiants, and lines represent upper limits (for O)




Fig. 6. Run of the c_1 index from Strömgren photometry with the Na abundance for stars in NGC6752. Filled squares are TO-stars, and open squares are subgiants

there is a clear anticorrelation between Mg and Al. Stars 1460 and 1481 (the two most Na-rich subgiants) are clearly Mg-poor, Al-rich. We underline here that while the Mg-Al anticorrelation is not obvious in panel a of Figure 7, we think that such an anticorrelation exists also for TO-star because the Al abundances show a very large star-to-star variation,




Fig. 7. Run of the abundances of Mg against that of Al for stars in NGC6752. The upper panel shows results for TO-stars; the lower panel those for subgiants. Note that Al abundances for TO-stars were obtained using the resonance doublet, that is affected by large departures from LTE. This causes the large offset of Al abundances obtained by the two groups of stars

correlated with Na abundance variations; we think the anticorrelation is less obvious in this case because we had to measure the resonance doublet of Al at 3944/61 Å, not an ideal abundance indicator as we mentioned above, and moreover Mg abundances have larger errors, comparable to the star-to-star variations seen among subgiants.

We are now preparing line lists to study CH and CN abundances. However, it seems clear that the strength of these bands is correlated with Na and O abundances. Also, there is a correlation with the Stromgren c_1 index, albeit the scatter is quite large. Finally, in a forthcoming paper we will present a full discussion of the Li abundances in these stars. Here we anticipate that Li abundances in NGC6752 stars seem also anticorrelated with the Na ones.

6. SUMMARY AND CONCLUSIONS

We have presented first results of the ESO Large Program 165.L-0263: we have used the UVES spectrograph on VLT2 to obtain high resolution ($R \gtrsim 40,000$) spectra for a quite large number of stars at the turn-off (14 stars between the two GCs) and the base of the subgiant branch (12 stars) in the globular clusters NGC6397 and NGC6752. Thanks to the efficiency and large spectral coverage of UVES we were able to obtain reliable *EWs*

for a number of lines of Fe, Li, O, Na, and other elements. The main results of this first analysis are:

(i) The $[\text{Fe}/\text{H}]$ value for NGC6397 is $[\text{Fe}/\text{H}] = -2.03 \pm 0.02 \pm 0.04$ (internal and systematic errors), less than derived from giants by Carretta & Gratton (1997) and Zinn & West (1984), but in agreement with the recent analysis of giants and subgiants by Castilho et al. (2000). The $[\text{Fe}/\text{H}]$ value for NGC6752 ($[\text{Fe}/\text{H}] = -1.42 \pm 0.02 \pm 0.04$) coincides with the value derived from giants by Carretta & Gratton (1997), and is slightly higher than the value quoted by Zinn & West (1984)

(ii) In both clusters, $[\text{Fe}/\text{H}]$ obtained for TO stars agrees perfectly (within a few percents) with that obtained for stars at the base of the giant branch; this is a constraint on the impact of diffusion in stellar models. The star-to-star scatter is extremely small: the r.m.s. scatter is only 0.04 dex (i.e. 10%) for NGC 6397. This seems a very homogeneous cluster as far as Fe abundances are considered.

(iii) For NGC6397, the O abundance is $[\text{O}/\text{Fe}] = +0.21 \pm 0.05$. This is a quite low value in comparison with those usually found in metal-poor stars, but it agrees well with previous determinations for red giants. The scatter of individual values is small, and none of the program stars seem to be oxygen-poor.

(iv) For NGC6752, there is a clear anticorrelation between Na and O among TO stars and subgiants (similar to that seen among giants in this and other clusters). Also, an anticorrelation is observed between Mg and Al, most clearly among subgiants, but likely existing also among TO-stars. Na-poor stars (i.e. stars with $[\text{Na}/\text{Fe}] < 0.2$) in NGC6752 have $[\text{O}/\text{Fe}] = +0.30 \pm 0.04$ (7 stars, r.m.s.=0.10 dex). Extremes in Al abundances differ by over 1 dex, while for Mg the star-to-star scatter is smaller. The large variations in Mg and Al abundances suggests that nearly half of Mg has been converted into Al in the most Al-rich stars (1481 and 1460); note that these stars are also the most Na-rich ones. Note that given the adopted sample selection criteria, extreme cases of Na-poor and Na-rich stars may be overrepresented among observed stars.

We think the present results are very difficult to be reconciled with deep mixing scenarios. To our knowledge, there are not appropriate calculations for main sequence stars. For giants, some calculations have been made by Langer, Hoffman & Sneden (1993) and Denissenkov & Tout (2000): they show that the temperature required for p-capture on ^{24}Mg to finally produce the Mg-Al anticorrelation is $\sim 6 \times 10^7$ K; even if Al is produced starting from the far less abundant ^{25}Mg isotope, the temperature required is $\gtrsim 4 \times 10^7$ K; finally, the temperature required for extensive O-burning and Na production by p-capture on ^{22}Ne is $\sim 3 \times 10^7$. All these values are much higher than expected even at the center of a globular cluster TO-star ($\sim 2 \times 10^7$ K). Admittedly, these computations assume densities about 20 times lower than expected at the center of TO-stars, and much shorter timescales than MS lifetimes. However complete mixing of MS stars is unacceptable for several other reasons (they would e.g. bring large amount of fresh H to

the center); furthermore, Li would be completely destroyed (while we see some Li even in O-depleted stars; paper in preparation). Hence, we think deep mixing scenarios cannot explain our results.

We are then forced to some primordial mechanism, like those proposed years ago by Cottrell & Da Costa (1981) and D’Antona, Gratton & Chieffi (1983). In both scenarios, the inhomogeneities are due to the mass lost by intermediate mass stars ($M = 4 - 5M_{\odot}$) during the Asymptotic Giant Branch (AGB) evolution and the planetary nebula expulsion: the two scenarios differ because Cottrell & Da Costa think of a prolonged star formation, with most recently formed stars having a different chemical composition from the first ones; while D’Antona et al. consider pollution of the outer layers of already formed stars by other objects in the cluster. Anyhow, in both scenarios the intracluster gas is heavily nucleary processed, due both to the occurrence of the third dredge-up from the helium buffer (Iben 1975) and by Hot Bottom Burning (HBB) at the basis of the convective envelopes of these massive AGB stars. Models for these evolutionary phases (Sackmann and Boothroyd 1992, Ventura et al. 2000) successfully explain the occurrence and evolution of the Lithium rich, Oxygen rich massive AGBs in the Magellanic Clouds (Smith et al. et al. 1995). Very recent models by Ventura et al. (2001) show that, in full stellar models computed for these intermediate mass stars at the low metallicities of Globular Clusters, the HBB temperature can reach values as large as 10^8K . At this temperature, the complete CNO cycle operates, depleting oxygen. At the same time, p-captures on ^{24}Mg and ^{20}Ne produce Al (see also Denissenkov et al. 1998) and Na. This model can then explain the anticorrelations O-Na and Mg-Al. In this context it is very interesting to note that no O-Na anticorrelation is seen in NGC6397. Also, Li abundances in this cluster follows the paradigm set by field metal-poor stars (Castilho et al. 2000). NGC6397 is a quite small cluster (mass $\lesssim 10^5 M_{\odot}$, from the integrated magnitude $M_V = -6.58$, Harris 1996, and a mass-to-light ratio of ~ 2 , typical for a globular cluster). On the other side, the O-Na anticorrelation is seen in the more massive cluster NGC6752 (mass $\sim 2 \cdot 10^5 M_{\odot}$, from the integrated magnitude $M_V = -7.68$, Harris 1996, and the same mass-to-light ratio used for NGC6397). A (cluster) mass threshold should be present in accretion scenarios (see e.g. Gratton 2001), and likely also in the prolonged star formation models. We plan to address thoroughly such problems in forthcoming papers.

Acknowledgements. This research has made use of the SIMBAD data base, operated at CDS, Strasbourg, France. We wish to thank V. Hill for help during the observations, and P. Bertelli for useful comments

References

- Alonso, A., Arribas, S., & Martinez-Roger, C. 1996, A&A, 313, 873
 Bell, R.A., & Dickens, R.J. 1980, ApJ, 242, 657

- Beyley, M.J. & Cottrell, P.L. 1987, MNRAS, 224, 633
- Boesgaard, A.M., Deliyannis, C.P., Stephens, A., & King, J.R. 1998, ApJ, 493, 206
- Bragaglia, A., Carretta, E., Gratton, R.G., Tosi, M., Bonanno, G., Bruno, P., Calì, A., Claudi, R., et al. 2001, AJ, in press
- Cannon, R.D., Croke, B.F.W., Bell, R.A., Hesser, J.E., & Stathakis, R.A. 1998, MNRAS, 298, 601
- Carretta, E. 1994, Ph.D. Thesis, Un. Padua
- Carretta, E., & Gratton, R.G. 1997, A&AS, 121, 95
- Carretta, E., Gratton, R.G., & Sneden, C. 2000, A&A, 356, 238
- Castelli, F., Gratton, R.G., & Kurucz, R.L. 1997, A&A, 318, 841
- Castilho, B.V., Pasquini, L., Allen, D.M., Barbuy, B., & Molaro, P., 2000, A&A 361, 92
- Cavallo, R.M., Sweigart, A.V., & Bell, R.A. 1996, ApJ, 464, L79
- Charbonnel, C. 1994, A&A, 282, 811
- Cohen, J.G. 1999a, AJ, 117, 2428
- Cohen, J.G. 1999b, AJ, 117, 2434
- Cottrell, P.L., & Da Costa, G.S. 1981, ApJL, 245, L79
- D’Antona, F., Gratton, R.G., & Chieffi, A. 1983, MSAIt, 54, 173
- Deliyannis, C.P., Boesgaard, A.M., & King, J.R. 1995, ApJ, 452, 12
- Denissenkov, P.A., & Denissenkova, S.N. 1990, SVAL, 16, 275
- Denissenkov, P.A., & Tout, C.A. 2000, MNRAS, 316, 395
- Denissenkov P.A., Da Costa, G.S., Norris, J.E., & Weiss, A. 1998, A&A, 333, 926
- D’Odorico, S., Cristiani, S., Dekker, H., Hill, V., Kaufer, A., Kim, & T. Primas, F. 2000, in Discoveries and Research Prospects from 8- to 10-Meter-Class Telescopes, J. Bergeron ed., SPIE Vol. 4005, p. 121
- François, P. 1986, A&A 160, 264
- Fuhrmann, K., Axer, M., & Gehren, T. 1993, A&A, 271, 451
- Fuhrmann, K., Axer, M., & Gehren, T. 1994, A&A, 285, 585
- Gratton, R.G., 2001, in The Chemical Evolution of the Galaxy, F. Matteucci and F. Giovannelli eds., Kluwer, Dordrecht, in press
- Gratton, R.G. & Sneden, C. 1988, A&A 204, 193
- Gratton, R.G., Carretta, E., Eriksson, K., & Gustafsson, B. 1999, A&A 350, 955
- Gratton, R.G., Sneden, C., Carretta, E., & Bragaglia, A. 2000, A&A, 354, 169
- Grundahl, F., Vandenberg, D.A., Stetson, P.B., Anderson, M.I., & Briley, M. 1999, in The Galactic Halo: from Globular Clusters to Field Stars (astro-ph/9909447)
- Harris, W.E. 1996, AJ, 112, 1487
- Idiart, T. & Thévenin, F. 1999, ApJ, 541, 207
- Iben, I.Jr. 1975, ApJ, 196, 525
- King, J., Stephens, A., & Boesgaard, A.M. 1998, AJ, 115, 666
- Kraft, R.P. 1994, PASP, 106, 553
- Kraft, R.P., Sneden, C., Smith, G.H., Shetrone, M.D., Langer, G.E., & Pilachowski, C.A. 1997, AJ, 113, 279
- Kurucz, R.L. 1995, CD-ROM 13, Smithsonian Astrophysical Observatory, Cambridge

- Langer, G.E., & Hoffman, R. 1995, *PASP*, 107, 1177
- Langer, G.E., Hoffman, R., & Sneden, C. 1993, *PASP*, 105, 301
- Minniti, D., Geisler, D., Peterson, R.C., & Claría, J.J. 1993, *ApJ*, 413, 548
- Minniti, D., Peterson, R.C., Geisler, D., & Claría, J.J. 1996, *ApJ*, 470, 953
- Norris, J.E. 1987, *ApJ*, 313, 65
- Norris, J.E., & Da Costa, G.S. 1995, *ApJL*, 441, L81
- Osborn, W. 1971 *Observatory*, 91, 223
- Pasquini, L., & Molaro, P. 1996, *A&A*, 307, 761
- Pasquini, L., & Molaro, P. 1997 *A&A*, 322, 109
- Sackmann, I.J., Boothroyd, A.I. 1992, *ApJ*, 392, L71
- Smith, G.H. 1987, *PASP*, 99, 67
- Smith, V.V., Plez, B., Lambert, D.L. 1995, *ApJ*, 441, 735
- Suntzeff, N.B., 1981, *ApJS*, 47, 1
- Suntzeff, N.B., & Smith, V. 1991, *ApJ*, 381, 160
- Sweigart, A.V., & Mengel, J.G. 1979, *ApJ*, 229, 624
- Ventura, P. D’Antona, F., Mazzitelli, I. 2000, *A&A* 363, 605
- Ventura, P. D’Antona, F., Mazzitelli, I., & Gratton, R. 2001, in preparation
- Zinn, R., & West, M.J. 1984, *ApJS*, 55, 45